



MANICALAND STATE UNIVERSITY OF APPLIED SCIENCES

FACULTY OF ENGINEERING

CHEMICAL AND PROCESSING ENGINEERING

PHYSICAL CHEMISTRY FOR ENGINEERS/ PHYSICAL CHEMISTRY

CODE: ENGT104/HCHE112

SESSIONAL EXAMINATIONS

OCTOBER 2021

LECTURER: MR M. MAPOSA

DURATION: 3 HOURS

INSTRUCTIONS

1. Answer ***all*** questions in section A and ***three*** questions in section B
2. Each question carries 25 marks
3. Total marks 100

ADDITIONAL MATERIAL

Periodic table

SECTION A

Answer all questions in this section

QUESTION 1

(a) State any two limitations of classical mechanics which necessitated the birth of quantum mechanics [2 marks]

(b) Explain the following terms i) quantum

ii) wave particle duality [4 marks]

(c) State first **three** postulates of quantum mechanics [3 marks]

(d) Write brief notes on

i) unimolecular reactions

ii) bimolecular reactions

iii) rate equation

iv) steady state approximation

v) mechanism of reaction [10 marks]

(e) Describe the instrumentation and applications of the following spectroscopic techniques

i. UV-Vis

ii. FTIR

iii. iii) Raman Spectroscopy [6 marks]

SECTION B

Answer **any three** questions in this section

QUESTION 2

In this question you may want to use the following constants:

1 eV = 1.602×10^{-19} J, $h = 6.63 \times 10^{-34}$ Js, mass of an electron = 9.11×10^{-31} kg

- a) During a lecture on photoelectric effect, the lecturer gave this statement as part of the introduction; “One of the factors that affect photoelectric emission is the threshold frequency of the metal” Explain the meaning of the terms
- photoelectric equation
 - Photoelectric emission
 - Threshold frequency [6 marks]
- b) Suggest any other factor besides threshold frequency which affect photoelectric emission [1 marks]
- c) i) Briefly explain the three laws of photoelectric emission
ii) Give any **four** applications of photoelectric emission [7 marks]
- d) The work function of zinc, silver and sodium are 4.31 eV, 4.73 eV and 2.28 eV respectively. Use the photoelectric equation to calculate
- Maximum kinetic energy of an electron ejected from silver by a 3.13×10^{15} hz [3 marks]
 - Maximum kinetic energy and velocity of the ejected electrons from zinc using 275 nm photon [3 marks]
 - The threshold frequency for the emission of photoelectrons from the surface of sodium metal, and hence the maximum wavelength of light that can cause photoemission. [5 marks]

QUESTION 3

- Sketch a graph to show how rate of reaction depends on concentration in a first order reaction. [3 marks]
- State the Arrhenius equation and explain each term present in the equation [4 marks]
- The rate constants for the reaction $A_2 + B_2 = 2AB$ were measured at five different temperatures. The results are shown in Table 2

Table 2

Experiment	Rate constant (k)/ s ⁻¹	Temperature/K
1	1.25 x 10 ⁻⁹	250
2	2.27 x 10 ⁻⁸	303
3	6.81 x 10 ⁻⁷	400
4	3.73 x 10 ⁻⁶	500
5	4.11 x 10 ⁻⁷	667

- i) Rewrite Arrhenius equation using natural logarithms
- ii) Produce a completed table of ln k and 1/T using the data from the table
- iii) Plot a graph of ln K against 1/T
- iv) Use the graph to deduce the activation energy of the reaction [13 marks]
- d) In another experiment the rate constant of the reaction $\text{H}_2 + \text{I}_2 = 2\text{HI}$ was measured at two different temperatures.

At **600 K**, $k = 2.15 \times 10^{-7} \text{ s}^{-1}$ At **700 K**, $k = 2.39 \times 10^{-6} \text{ s}^{-1}$

Calculate the activation energy of the reaction [5 marks]

Question 4

- a) What do you understand by the term complex reaction? [2 marks]
- b) Compare and contrast the Linderman and the RRKM theories [6 marks]
- c) Describe
- Chain reactions
 - Reaction intermediate
 - Elementary step of a mechanism
 - Adiabatic modes of vibration
 - Active modes of vibration [10 marks]
- d) Steady state approximation can be used to deduce the rate law from elementary steps. Given that



$A^* + M \xrightarrow{k_1} A + M$ Deactivation of A^* via collisions

$A^* \xrightarrow{k_2} P$ Spontaneous decomposition of A^*

(where M is any particle which can be an inert particle, a molecule of the product P or another particle of A), use steady state approximation to deduce that

i) $[A^*] = k_1[A][M] / k_{-1}[M] + k_2$

ii) $dP/dt = k_2 k_1[A][M] / k_{-1}[M] + k_2$ [7 marks]

QUESTION 5

a) State Beer's law [1 mark]

b) Describe the concept of molecular spectroscopy [3 marks]

c) An aqueous solution of substance X was analysed using UV-VIS at a wavelength of 275 nm at which molar absorptivity is $\epsilon = 8400 \text{ M}^{-1}\text{cm}^{-1}$. The pathlength was 1 cm. the absorbance (A_{275}) was found to be 70. What was the concentration of X in the sample? [5 marks]

d) During the analysis of dye concentration in a sample, a set of five standards were analysed at 500 nm to produce a calibration curve before running the sample. The equation of the line of best fit was found to be $y = 1.65x - 0.05$ where y is the absorbance, x is the concentration. If the sample gave an absorbance reading of 0.85, what will be the concentration of the dye in the sample? [4 marks]

e) Calculate the number of degrees of freedom for the following molecules

i) CO_2

ii) SO_2

iii) CH_4

iv) HCN [12 marks]

END OF EXAM