



MANICALAND STATE UNIVERSITY OF APPLIED SCIENCES

FACULTY OF ENGINEERING, SCIENCE AND TECHNOLOGY

DEPARTMENT: CHEMICAL AND PROCESSING ENGINEERING

MODULE: SEPARATION PROCESSES 1

CODE: CHEP 315

SESSIONAL EXAMINATIONS

APRIL 2023

DURATION: 3 HOURS

EXAMINER: MISS N.T. MADZIWA

INSTRUCTIONS

1. Answer *ALL* questions
2. Start a new question on a fresh page
3. Total marks 100
4. Formulae sheet is given at the end of the paper.

Additional material(s): Calculator

QUESTION 1

- a) Explain what is meant by *separation processes* in relation to chemical engineering operations. [2]
- b) State the applications of these **two** membrane processes:
- i. *Ultrafiltration*
 - ii. *Nanofiltration* [4]
- c) With the aid of well-labelled diagrams, explain the principle of reverse osmosis. [8]
- d) Explain these **two** types of membrane modules using well-labelled diagrams,.
- i. *Spiral-Wound Module*
 - ii. *Tubular Module* [6]

QUESTION 2

- a) Explain the process of electrodialysis using Na_2SO_4 as an electrolyte solution using an appropriate diagram [8]
- b) Describe the parameters influencing the efficiency of electrodialysis. [6]
- c) Explain the meaning of these terms as used in thermodynamics
- i. *Surroundings*
 - ii. *Entropy*
 - iii. *Closed system*
 - iv. *Gibbs energy* [4]
- d) Explain flow work and shaft work. [2]

QUESTION 3

- a) With the aid of diagrams, explain the process of crystallization and its applications in separation processes. [6]
- b) Explain the types of flow patterns as used in process mixing using appropriate diagrams [6]

c) With the aid of diagrams, explain these **two** mechanisms of equipment for solid mixing.

i. *Paddle mixer*

ii. *Tumbler mixer*

[8]

QUESTION 4

a) What is the yield of sodium acetate crystals ($\text{CH}_3\text{COONa}\cdot 3\text{H}_2\text{O}$) obtainable from a vacuum crystalliser operating at 1.33 kN/m^2 when it is supplied with 0.56 kg/s of a 40 per cent aqueous solution of the salt at 353 K ? The boiling point elevation of the solution is 11.5 deg K . [6]

Data:

Heat of crystallisation, $q = 144 \text{ kJ/kg}$ trihydrate

Heat capacity of the solution, $C_p = 3.5 \text{ kJ/kg deg K}$

Latent heat of water at 1.33 kN/m^2 , $\lambda = 2.46 \text{ MJ/kg}$

Boiling point of water at $1.33 \text{ kN/m}^2 = 290.7 \text{ K}$

Solubility of sodium acetate at 290.7 K , $c_2 = 0.539 \text{ kg/kg}$ water.

b) Describe the two types of primary nucleation. [4]

c) Explain these **two** types of the mechanisms of blending.

i. *Convective Mixing (Macro mixing)*

ii. *Diffusion mixing (Micro mixing)*

[6]

d) State the limitations of reverse osmosis (RO). [4]

QUESTION 5

a) (i) How much energy is available for sustaining muscular and nervous activity from the combustion of 1.00 mol of glucose molecules under standard conditions at $37 \text{ }^\circ\text{C}$ (blood temperature)? [2]

- (ii) Given the standard entropy of reaction as $+182.4 \text{ JK}^{-1} \text{ mol}^{-1}$. Outline the steps taken to calculate the energy available. [2]
- b) Given that the solubility of sodium sulphate at 283 K is 9 kg anhydrous salt/100 kg water and the deposited crystals will consist of the deca-hydrate (molecular mass = 322 kg/kmol). It may be assumed that 2 per cent of the water will be lost by evaporation during cooling). Calculate is the theoretical yield of crystals which may be obtained by cooling a solution containing 1500 kg of sodium sulphate (molecular mass =142 kg/kmol) in 5000 kg water to 283 K? [6]
- c) State the factors affecting mixing. [4]
- d) State the advantages of using tumbling mixers in solid mixing. [4]
- e) What is blending? [2]

END OF EXAMINATION

LIST OF FORMULAE

Crystal yield initial solvent balance: $w_1 = w_2 + y \frac{R-1}{R} + w_1 E$

Crystal yield solute balance: $w_1 c_1 = w_2 c_2 + y/R$

Yield for aqueous solutions: $y = R w_1 \frac{c_1 - c_2 (1-E)}{1 - c_2 (R-1)}$

Quantity from heat balance: $E = \frac{qR(c_1 - c_2) + C_p(T_1 - T_2)(1 + c_1)[1 - c_2(R-1)]}{\lambda[1 - c_2(R-1)] - qRc_2}$

Power: $\frac{\text{Power}_{gassed}}{\text{Power}_{ungassed}} = 0.1 \left(\frac{Q}{NV_L} \right)^{-0.25} \left(\frac{N^2 d^4}{gBV_L^{2/3}} \right)^{-0.20}$